

APPLICATION SER. NO. 10/722,679 CLAIMS

Please amend the claims of Application Ser. No. 10/722,679 as follows:

1. (Four Times Amended) A sweetener composition, comprising a mixture of:

(a) N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester; and

(b) Acesulfame K,

wherein said Acesulfame K is present in said mixture in an amount of 50 to 97% by weight

based on the total weight of said

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said Acesulfame

K and wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester

comprises a C-type crystal.

12. (Four Times Amended) A method for preparing a drink composition, comprising dissolving:

(A) a mixture, comprising:

(a) N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester; and

(b) Acesulfame K; [[and]]

in

(B) a potable liquid,

wherein said Acesulfame K is present in said mixture in an amount of 50 to 97% by weight

based on the total weight of said

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said Acesulfame

K and wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester

comprises a C-type crystal.

13. (Twice Amended): A method for preparing a sweetener composition, comprising:

(1) drying A-type crystals of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester, to obtain C-type crystals of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester; and

(2) mixing said C-type crystals of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester with Acesulfame K, to obtain a mixture,

wherein said Acesulfame K is present in said mixture in an amount of 50 to 97% by weight based on the total weight of said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said Acesulfame K, and wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester comprises a C-type crystal having a water content of less than 3% by weight.

14. (Three Times Amended): A method for producing a sweetener, comprising:

(1) mixing N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester with Acesulfame K, to obtain a mixture, wherein said Acesulfame K is present in said mixture in an amount of 50 to 97% by weight based on the total weight of said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said Acesulfame K and wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester comprises a C-type crystal.

16. (Three Times Amended): A method for improving the dissolution rate of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester, comprising:

(1) mixing said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester with Acesulfame K, prior to dissolving said

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester, to obtain a mixture,

wherein said Acesulfame K is present in said mixture in an amount of 50 to 97% by weight based on the total weight of said

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said Acesulfame K and wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester comprises a C-type crystal.

APPLICATION SER. NO. 90/007,160 CLAIMS

Please amend the claims of Application Ser. No. 90/007,160 as follows:

13. (Twice Amended): A method for preparing a sweetener composition, comprising:

(1) drying A-type crystals of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester, to obtain C-type crystals of

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester; and

(2) mixing said C-type crystals of
N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester with Acesulfame K,
to obtain a mixture,

wherein said Acesulfame K is present in said mixture in an amount of 50 to 97% by
weight based on the total weight of said

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said Acesulfame
K, and wherein said

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester comprises a C-type
crystal having a water content of less than 3% by weight.